

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
18 April 2002 (18.04.2002)

PCT

(10) International Publication Number
WO 02/30901 A1(51) International Patent Classification⁷: C07D 213/73,
471/14, A61K 31/5513, A61P 31/18

(21) International Application Number: PCT/US01/42143

(22) International Filing Date:
14 September 2001 (14.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/239,300 10 October 2000 (10.10.2000) US
60/291,841 17 May 2001 (17.05.2001) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- entirely in electronic form (except for this front page) and available upon request from the International Bureau

(71) Applicant: BOEHRINGER INGELHEIM CHEMICALS, INC. [US/US]; 2820 North Normandy Drive, Petersburg, VA 23113 (US).

(72) Inventor: GUPTON, Bernard, Franklin; 2601 Helmsley Court, Midlothian, VA 23113 (US).

(74) Agents: RAYMOND, Robert, P. et al.; Boehringer Ingelheim Corporation, 900 Ridgebury Road, P.O. Box 368, Ridgefield, CT 06877-0368 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 02/30901 A1

(54) Title: PROCESS FOR MAKING 3-AMINO-2-CHLORO-4-METHYLPYRIDINE

(57) Abstract: An improved process for the preparation of 2-chloro-3-amino-4-methylpyridine (CAPIC) which comprises the following steps: (a) reacting acetylacetaldehyde dimethyl acetal with malononitrile to yield a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene, (b) treating the mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene so produced with a strong acid and water, to yield 3-cyano-4-methyl-2-pyridone, (c) treating the 3-cyano-4-methyl-2-pyridone with a strong chlorinating agent, to yield 3-cyano-2-chloro-4-methylpyridine, (d) treating the 3-cyano-2-chloro-4-methylpyridine with a strong acid and water, to yield 2-chloro-3-amido-4-methylpyridine, and (e) treating the 2-chloro-3-amido-4-methylpyridine with a strong base and a halide, to yield 2-chloro-3-amino-4-methylpyridine.

PROCESS FOR MAKING 3-AMINO-2-CHLORO-4-METHYLPYRIDINE


5

Field of the Invention

10 The invention relates to an improved process for making 3-amino-2-chloro-4-methylpyridine, also known as CAPIC.

Background of the Invention

CAPIC is a key intermediate in the production of nevirapine, a non-nucleosidic reverse
15 transcriptase inhibitor that has been established to be clinically useful for the treatment of infection by HIV-1.




(nevirapine)

20

Syntheses of nevirapine from CAPIC have been described by Hargrave et al., in *J. Med. Chem.* 34, 2231 (1991) and U.S. Patent 5,366,972, and by Schneider et al., in U.S. Patent 5,569,760.

25 Several processes for preparing CAPIC have been described in the literature.


It is believed that the earliest synthesis of CAPIC, depicted below in Scheme 1, is that of Chapman et al. (*J. Chem Soc. Perkin Trans. 1*, (1980), 2398-2404).



Scheme 1

5

A closely related synthesis for CAPIC, depicted below in Scheme 2, has been described by Hargrave et al. (U.S. Patent 5,366,972).




Scheme 2

10

As reported by Grozinger et al. (*J. Heterocyclic Chem.*, **32**, 259 (1995)), CAPIC has been synthesized in small laboratory batches by nitrating the readily available 2-amino-4-picoline or 2-hydroxy-4-picoline, as depicted below in Scheme 3. This procedure suffers from non-selective nitration at positions 3 and 5, as well as thermo-chemical hazards and potential for "run-away" when carried out in large quantities.


15




Scheme 3

The drawbacks of the nitration-based process lead Grozinger to develop the two synthetic routes, which start from ethylacetoacetone and cyanacetamide, that are described in U.S.

Patent Nos. 5,668,287 and 5,200,522, and are depicted below in Schemes 4 and 5. Both of the latter two synthetic routes require the dichlorination of the intermediate 2,6-dihydroxy-4-methyl-3-pyridinecarbonitrile, at positions 2 and 6, subsequent de-chlorination and finally selective re-chlorination, using chlorine gas, at position 2. The di-chlorination and dehalogenation, as well as the selective monochlorination at position 2 require special manufacturing equipment that is expensive and may not be readily available.




Scheme 4



Scheme 5


Schneider (U.S. Patent 5,686,618) has provided an alternative means for mono-chlorinating 3-amino-4-methylpyridine at position 2, using H_2O_2 in HCl , instead of
5 chlorine gas.

Yet another synthesis, depicted below in Scheme 6, comprising the steps of chlorination of ethyl cyanoacetate, Michael addition with crotonaldehyde, cyclization, conversion to the amide and finally reduction to the amine has been described by Zhang et al. (*Tetrahedron*
10 51(48), 13177-13184 (1995)), who report that while the desired product was obtained, the Michael addition was slow and the cyclization low-yielding.




Scheme 6

A synthesis beginning with 2-chloro-3-aminopyridine has been disclosed by Nummy (U.S. Patent 5,654,429). This is depicted below in Scheme 7.




Scheme 7

International Application WO 00/43365 describes the process for preparing CAPIC that is depicted below in Scheme 8.




Scheme 8

A closely related alternative, depicted below in Scheme 9, is disclosed in International Application PCT/US00/00261 .




Scheme 9

- 5 Baldwin et al. (*J. Org. Chem.*, **43**, 2529 (1978)), reported a method for the preparation of 2-halonicotinic acid derivatives using β,γ -unsaturated aldehyde equivalents as shown in Scheme 10. The procedure involves the initial Knoevenagel condensation of an aldehyde or ketone with ethyl cyanoacetate or malonitrile followed by reaction with DMF acetal. The cyclization of the β,γ -unsaturated aldehyde equivalent is carried out by treatment with
- 10 HBr-acetic acid to give the 2-bromo adduct directly. Reaction yields with DMF acetal were in the 5 to 60% range, depending on the nature of the alkyl substituents. In several cases, the reaction of DMF acetal with the Knoevenagel adduct led to dimer formation. Overall yields for the two step process varied from 3 to 35% depending on the nature of the substituents.



Scheme 10

Baldwin et al., *supra.*, also used mixtures of acetals and enol ethers as β,γ-unsaturated aldehyde equivalents to overcome the limitations associated with the use of DMF acetal in the preparation of 2-bromonicotinic acid derivatives (Scheme 11). Yields also tended to vary with this approach depending on the substitution pattern of Knoevenagel adduct. Yields for the cyclization step ranged from 29 to 74% while overall yields for the two step process ranged from 15 to 40%.



Scheme 11

10

In summary, the methods that have been developed to date for the preparation of CAPIC and other related 4-alkylnicotinic acid derivatives suffer from excessive complexity,


inefficiency and/or lack of regiocontrol. Of the approaches considered, Baldwin's use of acetal/enol ether systems addresses the regioselectivity issues most effectively

5 Description of the Invention

In its most general aspect, the invention comprises an improved process for the preparation of 2-chloro-3-amino-4-methylpyridine (CAPIC) which comprises the following steps:


(a) reacting acetylacetaldehyde dimethyl acetal

10




with malononitrile

15




to yield a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene

20




(b) treating the mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene so produced with a strong acid and water, to yield 3-cyano-4-methyl-2-pyridone




5

(c) treating the 3-cyano-4-methyl-2-pyridone so produced with a strong chlorinating agent, to yield 3-cyano-2-chloro-4-methylpyridine



10


(d) treating the 3-cyano-2-chloro-4-methylpyridine produced in the preceding step with a strong acid and water, to yield 2-chloro-3-amido-4-methylpyridine



15

(e) treating the 2-chloro-3-amido-4-methylpyridine produced in the preceding step with a strong base and a halide, to yield 2-chloro-3-amino-4-methylpyridine.

20 This general method is depicted below in Scheme 12.

**Scheme 12**

The intermediates 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene are believed to be novel, and constitute an aspect of the invention.

5

The first step of the above-described process is a Knoevenagel condensation. This is carried out in an organic solvent at a temperature in the range between 0 and 50°C. Acceptable solvents are, for example, aromatic hydrocarbons such as benzene, toluene or alkanols such as methanol, ethanol, propanols and other higher molecular weight alcohols. The preferred solvents are toluene and methanol. The preferred reaction temperature is 15 to 25 °C. The condensation is run in the presence of a small quantity of an ammonium salt catalyst. Preferred catalysts are heterocyclic ammonium salts. Most preferred is piperidinium acetate.

10

The product of the Knoevenagel condensation is a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene. These two compounds need not be separated, as they both are converted in the next step, which is an acid-catalyzed cyclization, to 3-cyano-4-methyl-2-pyridone. It is possible to conduct the cyclization simply by acidifying the initial reaction mixture, without isolation of the mixture of intermediates. However, it is preferred to first isolate the two intermediates from the other components of the crude Knoevenagel reaction mixture. This is conveniently accomplished by washing the reaction mixture emanating from the condensation with water, to remove the basic catalyst, followed by evaporation, to remove the solvent (toluene or methanol).

While the crude reaction mixture can be carried forward to the next step after catalyst removal and concentration, it has been discovered that this crude product contains undesired byproducts that tend to be carried forward into subsequent reaction steps and reduce yields. Therefore it is desirable to remove these byproducts once the catalyst removal and concentration have been completed. The removal of these byproducts can be performed by simple distillation techniques, preferably by continuous short path distillation such as wiped or thin film evaporation. This technique is particularly effective for the removal of high boiling materials, which appears to be sufficient to obtain the observed yield improvements.

The acid-catalyzed cyclization is performed with a strong acid, such as, for example, concentrated sulfuric acid. As the reaction is highly exothermic, it is best to introduce the mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene into the acid slowly, and with stirring, so that the temperature of the mixture does not rise above about 50°C. After evolution of heat has substantially ceased, the reaction mixture is heated to between about 30 and 50°C, preferably 50 °C, and held at that temperature for between about 1 and 3 hours, preferably about 1.5 hours, to complete the reaction. The reaction mixture is cooled to ambient temperature, water is added, and the intermediate product, 3-cyano-4-methyl-2-pyridone, is filtered off, washed with water and dried.

Next, the 3-cyano-4-methyl-2-pyridone so produced is treated with a strong chlorinating agent. Suitable chlorinating agents are SOCl_2 , POCl_3 and PCl_5 . It is preferred to use POCl_3 (10 parts) and PCl_5 (1 part). The reaction mixture is heated to reflux (approximately 5 115 °C) and held under this condition for about two hours, or until the chlorination is essentially complete. The chlorinating agent is removed. For example, excess POCl_3 may be removed by distillation. The reaction mixture is then cooled and water is added. The 2-chloro-3-cyano-4-methylpyridine is filtered from the aqueous mixture. The aqueous filtrate is extracted with an inert organic solvent such as chloroform, methylene chloride, 10 with methylene chloride being preferred in order to recover the residual 2-chloro-3-cyano-4-methylpyridine.

The 3-cyano intermediate is next converted to a 3-amido compound. This is accomplished by treating the cyano intermediate with a concentrated, aqueous strong acid, such as 15 sulfuric acid. This is preferably done with stirring, at a temperature between about 70 and 110°C, preferably at about 90 °C. The mixture was heated to between about 80 and 120°C, preferably about 100 °C and held at that temperature for three hours, or until further reaction ceases. The reaction mixture is then cooled to between about 70 and 110°C, preferably about 90°C, and water is added. The mixture is then cooled to between about 0 20 and 20°C, preferably about 10 °C and held at that temperature for about one hour. The solid product, 2-chloro-3-amido-4-methylpyridine, is isolated from the reaction mixture by filtration, washed with water and dried.


In the final step of the process, the 3-amido intermediate is converted to the 3-amino final 25 product by adding it to a mixture of a strong base and a halogen. The base may be aqueous sodium carbonate or sodium hydroxide, preferably sodium hydroxide. The halogen may be chlorine or bromine, preferably bromine. The resulting reaction mixture is heated to between about 10 and 30°C, preferably to about 22 °C. Water is then added to the reaction mixture followed by heating to between 60 and 80°C, preferably to about 70 °C, for one 30 additional hour. The reaction mixture is cooled to ambient temperature and extracted with an inert organic solvent, such as chloroform or methylene chloride, preferably methylene

chloride. The organic solvent is removed by evaporation, to yield 2-chloro-3-amino-4-methylpyridine.

Example 1, below, describes a specific preparation of CAPIC that was carried out in accordance with the general method of the invention.

Example 1

- a) Preparation of a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene (Knovenagel Condensation)



Piperidinium acetate (5.42 g, 0.037 moles) was added to a stirred solution of toluene (150 ml) and acetylacetaldehyde dimethyl acetal (49.3 g, 0.373 moles). Malononitrile (24.65 g, 0.373 moles) was then added to the reaction mixture over a 20-minute period. The mixture was allowed to stir for 24 hours at room temperature. The reaction mixture was then washed with 50 mL of water followed by rotary evaporation to give 67.14 g of a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetatal (1) and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene (2).

mp: not isolated

Yield: 71.9% (1) and 21.6% (2)


¹H NMR: s 4.55 (m, 1H), 3.4 (s, 6H), 2.9 (d, 2H), 2.25 (s, 3H)

IR (KBr): 3059, 2219, 1629, 1547, 1359, 1248, 1143, 981 cm⁻¹

Mass spectrum m/z: (1) m/z 179, 149, 121, 75 (2) 148, 133, 119, 91, 83, 78

Purity: crude mixture

(b) Preparation of 3-cyano-4-methyl-2-pyridone
(Cyclization)



5

The mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal (1) and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene (2) produced by the preceding step was added to a stirred solution of concentrated sulfuric acid (109.8 g, 1.12 moles) at a rate so that the reaction contents did not exceed 30 °C. The reaction mixture was then heated to 50 °C and held at
 10 that temperature for 1.5 hours. The reaction mixture was cooled to ambient temperature followed by the addition of water (150 mL). The product was filtered from the reaction mixture, washed with water (20 mL) and dried in a vacuum oven at 70 °C and full vacuum for 12 hours to give 40.7 g of 3-cyano-4-methyl-2-pyridone (3).
 mp: 230-232 °C

15 Yield: 81.4%

^1H NMR: 6.95 (d, $J = 2\text{ Hz}$, 1H), 6.29 (d, $J = 2\text{ Hz}$, 1H), 2.35 (s, 3H); ^{13}C NMR 162.0, 160.4, 140.0, 116.1, 108.1, 103.4, 21.1


IR (KBr): 3144, 3097, 2979, 2833, 2227, 1652, 1635, 1616, 1539, 1484, 1242, 1218, 1173, 819 cm^{-1}

20 Mass spectrum m/z : 134, 105, 78, 63, 51

Purity: 98.6 % by HPLC analysis (peak area basis)

(c) Preparation of 2-chloro-3-cyano-4-methylpyridine
(Chlorination)

25



The product of the preceding step, 4-methyl-3-cyano-2-pyridone (40.7 g, 0.304 moles) was added to a stirred solution of POCl₃ (140 g, 0.912 moles) and PCl₅ (19.0 g, 0.091 moles).

5 The reaction was heated to reflux (approximately 115 °C) and held under those conditions for two hours. Excess POCL₃ was removed by distillation. The reaction mixture was then cooled followed by the addition of water (30 mL) water. The aqueous mixture was extracted with 30 mL methylene chloride. The extracts were concentrated and 43.2 g was recovered as a tan solid, which was determined to be 2-chloro-3-cyano-4-methylpyridine (4).

mp: 102-104 °C

Yield: 82.7 %


¹H NMR: 8.03, (d, J= 2 Hz, 1H), 7.6 (d, J= 2 Hz, 1H), 2.5 (s, 3H); ¹³C NMR 156.1, 152.6, 151.8, 124.9, 114.4, 111.8, 20.6

15 Mass spectrum m/z: 155, 154, 152, 117, 116, 90, 89, 76, 64, 63, 62

FTIR (KBr): 3144, 2979, 2834, 2228, 1653, 1616, 1540, 1484, 1242, 1218, 1173, 819, 607 cm^{-1}

Purity: 98.2% by HPLC analysis (peak area basis)

20 (d) Preparation of 2-chloro-3-amido-4-methylpyridine
(Addition of water)



The product of the preceding step, 2-chloro-3-cyano-4-methylpyridine (34.14 g, 0.224 moles) was added to a stirred solution of concentrated sulfuric acid (65.87 g, 0.672 moles) at 90 °C. The mixture was then heated to 100 °C and held at that temperature for three hours. The reaction was then cooled to 90 °C followed by the addition of water (121 mL). The mixture was cooled to 10 °C and held at that temperature for one hour. The solid product was isolated from the reaction mixture by filtration, washed with water (16 mL) and dried at 70 °C under full vacuum to give 34.7 g of 2-chloro-3-amido-4-methylpyridine (5).

mp: 178-179 °C

Yield: (90.9%)


¹H NMR: 8.28 (d, J= 8 Hz, 1H), 8.09 (s, 1H), 7.84 (s, 1H), 7.34 (d, J= 8 Hz, 1H), 2.31 (s, 3H)

¹³C NMR: 166.4, 1149.0, 1437.8, 146.2, 136.0, 125.0, 18.9

Mass spectrum m/z: 155, 154, 152, 116, 91, 90, 89, 88, 76, 75, 64, 63, 62, 52, 51

FTIR (KBr): 3407, 3303, 3194, 3116, 3069, 3031, 2962, 2847, 1664, 1539, 1475, 1458, 1380, 1174, 1145, 798, 595 cm⁻¹.

(e) Preparation of 2-chloro-3-amino-4-methylpyridine



Bromine (34.7 g, 0.204 moles) was added to a stirred solution of sodium hydroxide (8.14 g, 0.224 moles) and water (205 mL) at 0 °C. The product of the preceding step, 2-chloro-3-amido-4-picoline (34.7 g, 0.204 moles) was then added to the reaction mixture followed by heating to 22 °C. An additional 66 mL water was then added to the reaction mixture followed by heating to 70 °C and stirring for one additional hour. The reaction mixture was cooled to ambient temperature and extracted with methylene chloride (60 mL). The methylene chloride was removed by rotary evaporation to give 24.8 g of 2-chloro-3-amino-4-methylpyridine (6).

mp: 69 °C

Yield: 85.2%

¹H NMR: 7.5 (d, J= 4.64Hz, 1H), 7.0 (d, J= 4.28 Hz, 1H), 5.25 (s, 2H), 2.1 (s, 3H)

¹³C NMR: 140.0, 136.2, 135.6, 131.9, 125.7, 19.0


IR (KBr): 3429, 3308, 3198, 1630, 1590, 1550, 1475, 1451, 1441, 1418, 1377, 1295, 1122, 860, 821, 656, 527 cm⁻¹

Mass spectrum m/z: 145, 144, 142, 107, 106, 105, 80, 79, 78, 62, 54, 53, 52

Purity: 87.6%, HPLC analysis (peak area basis).

Example 2, below, describes a specific preparation of CAPIC that was carried out in accordance with the preferred method of the invention in which the Knoevenagel condensation reaction product is purified by short path distillation.


- 5 a) Preparation of a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene followed by short path distillation (Knoevenagel Condensation)



- 10 Piperidinium acetate (5.42 g, 0.037 moles) was added to a stirred solution of toluene (150 ml) and acetylacetaldehyde dimethyl acetal (49.3 g, 0.373 moles). Malononitrile (24.65 g, 0.373 moles) was then added to the reaction mixture over a 20-minute period. The mixture was allowed to stir for 24 hours at room temperature. The reaction mixture was then washed with 50 mL of water followed by rotary evaporation to give 67.14 g of a mixture of
- 15 4,4-dicyano-3-methyl-3-butenal dimethyl acetatal (1) and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene (2). The crude reaction product was then distilled on a 0.1 square meter stainless steel wiped film evaporator at 109 °C, 0.1 mm Hg and 240 RPM to give 60.1 kg of a yellow liquid

- 20 Yield: 85.0% (1) and (2)
- ¹H NMR: s 4.55 (m, 1H), 3.4 (s, 6H), 2.9 (d, 2H), 2.25 (s, 3H)
- IR (KBr): 3059, 2219, 1629, 1547, 1359, 1248, 1143, 981 cm⁻¹
- Mass spectrum m/z: (1) m/z 179, 149, 121, 75 (2) 148, 133, 119, 91, 83, 78
- Purity: 94.0% (1) and (2)

(b) Preparation of 3-cyano-4-methyl-2-pyridone
(Cyclization)



5

The mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal (1) and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene (2) produced by the preceding step was added to a stirred solution of concentrated sulfuric acid (109.8 g, 1.12 moles) at a rate so that the reaction contents did not exceed 30 °C. The reaction mixture was then heated to 50 °C and held at that temperature for 1.5 hours. The reaction mixture was cooled to ambient temperature followed by the addition of water (150 mL). The product was filtered from the reaction mixture, washed with water (20 mL) and dried in a vacuum oven at 70 °C and full vacuum for 12 hours to give 40.7 g of 3-cyano-4-methyl-2-pyridone (3).

mp: 230-232 °C

Yield: 90.6 %

^1H NMR: 6.95 (d, $J=2\text{Hz}$, 1H), 6.29 (d, $J=2\text{Hz}$, 1H), 2.35(s, 3H); ^{13}C NMR 162.0, 160.4, 140.0, 116.1, 108.1, 103.4, 21.1


IR (KBr): 3144, 3097, 2979, 2833, 2227, 1652, 1635, 1616, 1539, 1484, 1242, 1218, 1173, 819 cm^{-1}

Mass spectrum m/z : 134, 105, 78, 63, 51

Purity: 98.8 % by HPLC analysis (peak area basis)

25

(c) Preparation of 2-chloro-3-cyano-4-methylpyridine
(Chlorination)



5

The product of the preceding step, 4-methyl-3-cyano-2-pyridone (40.7 g, 0.304 moles) was added to a stirred solution of POCl_3 (140 g, 0.912 moles) and PCl_5 (19.0 g, 0.091 moles). The reaction was heated to reflux (approximately 115 °C) and held under those conditions for two hours. Excess POCl_3 was removed by distillation. The reaction mixture was then cooled followed by the addition of water (30 mL) water. The aqueous mixture was extracted with 30 mL methylene chloride. The extracts were concentrated and 43.2 g was recovered as a tan solid, which was determined to be 2-chloro-3-cyano-4-methylpyridine (4).

10

mp: 102-104 °C

15

Yield: 98.2 %

^1H NMR: 8.03 (d, $J = 2$ Hz, 1H), 7.6 (d, $J = 2$ Hz, 1H), 2.5 (s, 3H); ^{13}C NMR 156.1, 152.6, 151.8, 124.9, 114.4, 111.8, 20.6

Mass spectrum m/z : 155, 154, 152, 117, 116, 90, 89, 76, 64, 63, 62

FTIR (KBr): 3144, 2979, 2834, 2228, 1653, 1616, 1540, 1484, 1242, 1218, 1173, 819, 607


20

cm^{-1}

Purity: 98.6% by HPLC analysis (peak area basis)

25

(d) Preparation of 2-chloro-3-amido-4-methylpyridine
(Addition of water)



5

The product of the preceding step, 2-chloro-3-cyano-4-methylpyridine (34.14 g, 0.224 moles) was added to a stirred solution of concentrated sulfuric acid (65.87 g, 0.672 moles) at 90 °C. The mixture was then heated to 100 °C and held at that temperature for three hours. The reaction was then cooled to 90 °C followed by the addition of water (121 mL).

10 The mixture was cooled to 10 °C and held at that temperature for one hour. The solid product was isolated from the reaction mixture by filtration, washed with water (16 mL) and dried at 70 °C under full vacuum to give 34.7 g of 2-chloro-3-amido-4-methylpyridine (5).

mp: 178-179 °C

15 Yield: 92.93%

¹H NMR: 8.28 (d, J= 8 Hz, 1H), 8.09 (s, 1H), 7.84 (s, 1H), 7.34 (d, J= 8 Hz, 1H), 2.31 (s, 3H)


¹³C NMR: 166.4, 1149.0, 1437.8, 146.2, 136.0, 125.0, 18.9

Mass spectrum m/z: 155, 154, 152, 116, 91, 90, 89, 88, 76, 75, 64, 63, 62, 52, 51

20 FTIR (KBr): 3407, 3303, 3194, 3116, 3069, 3031, 2962, 2847, 1664, 1539, 1475, 1458, 1380, 1174, 1145, 798, 595 cm⁻¹.

25

(e) Preparation of 2-chloro-3-amino-4-methylpyridine



5

Bromine (34.7 g, 0.204 moles) was added to a stirred solution of sodium hydroxide (8.14 g, 0.224 moles) and water (205 mL) at 0 °C. The product of the preceding step, 2-chloro-3-amido-4-picoline (34.7 g, 0.204 moles) was then added to the reaction mixture followed by heating to 22 °C. An additional 66 mL water was then added to the reaction mixture followed by heating to 70 °C and stirring for one additional hour. The reaction mixture was cooled to ambient temperature and extracted with methylene chloride (60 mL). The methylene chloride was removed by rotary evaporation to give 24.8 g of 2-chloro-3-amino-4-methylpyridine (6).

mp: 69 °C


15 Yield: 93.1 %

 ^1H NMR: 7.5 (d, $J=4.64\text{Hz}$, 1H), 7.0 (d, $J=4.28\text{ Hz}$, 1H), 5.25 (s, 2H), 2.1 (s, 3H) ^{13}C NMR: 140.0, 136.2, 135.6, 131.9, 125.7, 19.0IR (KBr): 3429, 3308, 3198, 1630, 1590, 1550, 1475, 1451, 1441, 1418, 1377, 1295, 1122, 860, 821, 656, 527 cm^{-1} 20 Mass spectrum m/z : 145, 144, 142, 107, 106, 105, 80, 79, 78, 62, 54, 53, 52

Purity: 100.2 %, HPLC analysis (peak area basis).

What is claimed is:

1. A process for preparing 2-chloro-3-amino-4-methylpyridine



5

which comprises the following steps:

- (a) reacting acetylacetaldehyde dimethyl acetal with malononitrile, in the presence of an ammonium salt catalyst, to yield a mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene,
- 10 (b) treating the mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene so produced with a strong acid and water, to yield 3-cyano-4-methyl-2-pyridone,
- (c) treating the 3-cyano-4-methyl-2-pyridone so produced with a strong chlorinating agent,
- 15 (d) treating the 3-cyano-2-chloro-4-methylpyridine produced in the preceding step with a strong acid and water, to yield 2-chloro-3-amido-4-methylpyridine, and
- (e) treating the 2-chloro-3-amido-4-methylpyridine produced in the preceding step with a strong base and a halide, to yield 2-chloro-3-amino-4-methylpyridine.

20

2. The process of claim 1 wherein the ammonium salt catalyst used in step (a) is piperidinium acetate.
3. The process of claim 1 wherein the acid used in step (b) is sulfuric acid.
- 25 4. The process of claim 1 wherein the chlorinating agent used in step (c) is a mixture of POCL₃ and PCl₅.

5. The process of claim 1 wherein the strong acid used in step (d) is sulfuric acid.

6. The process of claim 1 wherein the base and halide used in step (e) are, respectively, NaOH and Br₂.

5

7. The process of claim 1 wherein the crude mixture of 4,4-dicyano-3-methyl-3-butenal dimethyl acetal and 1,1-dicyano-4-methoxy-2-methyl-1,3-butadiene produced in step (a) is washed with water to remove catalyst, followed by evaporation to remove solvent, and the remaining crude mixture is then subjected to distillation, to remove high boiling reaction byproducts, and then the remaining process steps (b) through (e) are carried out.

10

8. The process of claim 7, wherein the distillation is continuous, short path distillation.

15

9. The process of claim 8 wherein the distillation is accomplished via wiped or thin film evaporation.

10. 4,4-Dicyano-3-methyl-3-butenal dimethyl acetal.

20

11. 1,1-Dicyano-4-methoxy-2-methyl-1,3-butadiene.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/42143

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D213/73 C07D471/14 A61K31/5513 A61P31/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 43365 A (BOEHRINGER INGELHEIM PHARMA) 27 July 2000 (2000-07-27) cited in the application the whole document	1-11
A	----- HARGRAVE K D ET AL: "NOVEL NON-NUCLEOSIDE INHIBITORS OF HIV-1 REVERSE TRANSCRIPTASE. 1. TRICYCLIC PYRIDOBENZO- AND DIPYRIDODIAZEPINONES" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 34, no. 7, July 1991 (1991-07), pages 2231-2241, XP002912882 ISSN: 0022-2623 cited in the application the whole document -----	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the International search

12 February 2002

Date of mailing of the International search report

22/02/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax. (+31-70) 340-3016

Authorized officer

Bosma, P

Information on patent family members

PCT/US 01/42143

Form PGT/SA/210 (patent family annex) (July 1992)

Schedule Service

Thank you for scheduling your appliance service online

Confirmation number: 020990299 | Last Name: wales

Use this number and name to check the status, change or cancel your service.

[Print this page for your records](#)

Important information about your service appointment

- **On the evening before** your scheduled repair appointment, GE will call at your primary phone number to confirm your appointment.
- **On the day of** your appointment, you will also receive a call from a GE service technician to confirm the appointment details and verify that someone over 18 is at the appliance location. If the technician cannot reach you at your primary phone number, the alternate phone number (if applicable) will be used.

In the event that GE cannot reach you after at least two attempts within the appointment time span, the service call will be canceled. To reschedule the appointment, call us at 1-800-GEcares (1-800-432-2737).

Service requested for:

GE Profile
Side By Side Refrigerator

Scheduled Date:

Thursday, June 15
Afternoon, 1:00 pm to 5:00 pm

Customer name and location of appliance:

William wales
1272 Helford Lane
Carmel, IN 46032
317-843-1845
317-796-7099
wwwales@aol.com

[Schedule service for another appliance](#)

Prefer to schedule service by phone? Call 1-800-GEcares (1-800-432-2737)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.